

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory

ZITHROPYX

Azithromycin Tablets USP 500 mg

COMPOSITION :

Each film coated tablet contains:
Azithromycin Dihydrate USP
eq. to Azithromycin Base 500 mg
Excipients Q.S.
Colour: Titanium Dioxide BP

INDICATIONS:

Respiratory Tract Infection, Urinary Tract Infection, Skin & Soft tissue Infections, Genital Infection, Uncomplicated chlamydial Infection, Otitis media.

CONTRAINDICATIONS:

Hypersensitivity to ZITHROPYX [Azithromycin] or any macrolide antibiotics.
Co-administration with ergot derivatives.

PRECAUTIONS:

Hepatic and renal impairment.
Pseudomembranous colitis.
Antacids with Ca and Mg.

USE IN PREGNANCY:

ZITHROPYX [Azithromycin] does cross the placenta and should be used during pregnancy only if adequate alternatives are not available.

DRUG INTERACTIONS:

Disopyramide: Displaced from its protein-bound sites.
Digoxin: Increases its absorption.
Midazolam: Increases its half-life.
Terfenadine: Serum levels increased leading to ventricular arrhythmias.

ADVERSE REACTIONS:

Nausea, Vomiting, Diarrhoea/flatulence, Epigastric distress, Dizziness/vertigo, Headache, Urticaria, Rash, Jaundice, Cardiac effects.

OVERDOSAGE:

Emesis, gastric lavage. Administer activated charcoal. General supportive & symptomatic treatment. Dialysis is of no help.

DOSAGE & ADMINISTRATION:

Adults: URTI/LRTI, skin & soft tissue infection: 500mg OD-3 days or 500 OD on day, 1 followed by 250mg OD - 4days.

UTI (chlamydial) - 1g single dose.

Children: 10mg/kg/d x 3days or 10mg/kg/d on day one followed by 5mg/kg/once daily for 5 days or as directed by the Physician.

STORAGE :

Store below 30°C. Protect from light and moisture.

Keep the medicine out of reach of children.

PRESENTATION :

3 Tablets

Marketed by:



PYXUS PHARMACEUTICALS PVT. LTD.

A/707, Mondeal Heights, Beside Novotel Hotel,
Nr. Iscon Square, S.G. Highway, Ahmedabad,
Gujarat- 380015 (India).